## Madagascar Medical Report – 2025

Category	Description	N Cases	Nature of illness/injury	Treatment and action taken	Cases/1000 person days
1	Death	0	N/A	N/A	0.00
2	Emergency evacuation with subsequent repatriation for medical attention	0	N/A	N/A	0.00
3	Non-Emergency evacuations	0	N/A	N/A	0.00
4	Hospital Treatment required in the local area	3	Cat bite and/or dog scratch	Began course of post-exposure rabies vaccinations at hospital in Nosy Be.	0.72
5	Treatment on site- serious illness/injury	0	N/A	N/A	0.00
6	Treatment on site- moderate serious illness/injury	28	Serious cases of D&V, more serious allergic reactions/rashes, cuts/abrasions with greater risk of infection/deeper or more extensive abrasions, more serious side effects from anti-malaria medication, serious cases of dehydration and/or heat exhaustion, more severe chest/abdominal pain.	Patients with more serious cases of D&V were treated by on-site medic. Electrolytes, paracetamol administered and where necessary a course of antibiotics given.  Any serious allergic reactions/rashes treated with antihistamines (oral or topical) and monitored for changes.  Any more serious cuts/abrasions were cleaned and dressed and antibiotics administered if necessary.  For more serious side effects from anti-malaria medication, an alternative prophylaxis was recommended and administered.  Any cases of dehydration and/or heat exhaustion were treated with fluids and electrolytes, patients were closely monitored for changes.	6.72
7	Treatment on site - minor illness/injury	201	Minor cuts and abrasions, minor gastrointestinal problems, muscular pain, common cold, nausea, diarrhoea and vomiting, insect bites/stings, blisters, constipation, splinters, joint pain, sore throat/cough, mild fever, travel sickness, nosebleeds, headaches, abdominal cramps, minor rashes, dental concerns, asthma attack, ear pain/infection, seasickness, blocked sinuses/congestion, sunburn, perforated eardrum	The majority of cases reported were minor gastrointestinal issues and cases of nausea, diarrhoea and vomiting. These were often treated with electrolytes, paracetamol, and plenty of fluids/rest was advised. Often resolved within 48 hours.  Bites, stings and rashes were treated with antihistamines (oral or topical) and cleaned where appropriate.  Minor cuts, abrasions, blisters, splinters were all cleaned and dressed were appropriate. Reviewed for signs of infection where appropriate.  Ear pain/infections treated with ear drops and congestion with decongestants and, where necessary, patients were kept out of the water and advised not to dive.  For perforated eardrum confirmed with examination. No diving/swimming advised. Regular analgesia given and advised to return if signs of infection. GP referral.  For sprained ankles/joint pain general RICE recommended. Analgesia offered and reassessment in UK recommended if pain persists.	48.26